

PubMed

Format: Abstract

Full text links

J Bone Miner Res. 2015 Oct;30(10):1905-13. doi: 10.1002/jbmr.2513. Epub 2015 May 10.



Cannabidiol, a Major Non-Psychotropic Cannabis Constituent Enhances Fracture Healing and Stimulates Lysyl Hydroxylase Activity in Osteoblasts.

Kogan NM¹, Melamed E¹, Wasserman E¹, Raphael B^{1,2}, Breuer A³, Stok KS⁴, Sondergaard R⁴, Escudero AV⁴, Baraghithy S¹, Attar-Namdar M¹, Friedlander-Barenboim S⁵, Mathavan N^{6,7}, Isaksson H^{6,7}, Mechoulam R³, Müller R⁴, Bajayo A¹, Gabet Y², Bab I¹.

Author information

Abstract

Cannabinoid ligands regulate bone mass, but skeletal effects of cannabis (marijuana and hashish) have not been reported. Bone fractures are highly prevalent, involving prolonged immobilization and discomfort. Here we report that the major non-psychoactive cannabis constituent, cannabidiol (CBD), enhances the biomechanical properties of healing rat mid-femoral fractures. The maximal load and work-to-failure, but not the stiffness, of femurs from rats given a mixture of CBD and $\Delta(9)$ -tetrahydrocannabinol (THC) for 8 weeks were markedly increased by CBD. This effect is not shared by THC (the psychoactive component of cannabis), but THC potentiates the CBD stimulated work-to-failure at 6 weeks postfracture followed by attenuation of the CBD effect at 8 weeks. Using micro-computed tomography (μ CT), the fracture callus size was transiently reduced by either CBD or THC 4 weeks after fracture but reached control level after 6 and 8 weeks. The callus material density was unaffected by CBD and/or THC. By contrast, CBD stimulated mRNA expression of Plod1 in primary osteoblast cultures, encoding an enzyme that catalyzes lysine hydroxylation, which is in turn involved in collagen crosslinking and stabilization. Using Fourier transform infrared (FTIR) spectroscopy we confirmed the increase in collagen crosslink ratio by CBD, which is likely to contribute to the improved biomechanical properties of the fracture callus. Taken together, these data show that CBD leads to improvement in fracture healing and demonstrate the critical mechanical role of collagen crosslinking enzymes.

© 2015 American Society for Bone and Mineral Research.

KEYWORDS: CANNABIDIOL; COLLAGEN CROSSLINKING; FRACTURE HEALING; FTIR; LYSYL HYDROXYLASE; μ CT

PMID: 25801536 DOI: [10.1002/jbmr.2513](https://doi.org/10.1002/jbmr.2513)

[Indexed for MEDLINE] [Free full text](#)