



Joints for joints: cannabinoids in the treatment of rheumatoid arthritis

Torsten Lowin, Matthias Schneider, and Georg Pongratz

Purpose of review

An increasing number of patients with rheumatoid arthritis (RA) are using cannabis to treat their symptoms, although systematic studies regarding efficacy in RA are lacking. Within this review we will give an overview on the overall effects of cannabinoids in inflammation and why they might be useful in the treatment of RA.

Recent findings

Peripherally, cannabinoids show anti-inflammatory effects by activating cannabinoid type 2 receptors (CB₂) which decrease cytokine production and immune cell mobilization. In contrast, cannabinoid type 1 receptor (CB₁) activation on immune cells is proinflammatory while CB₁ antagonism provides anti-inflammatory effects by increasing β_2 -adrenergic signaling in the joint and secondary lymphoid organs. In addition, the nonpsychotropic cannabinoid, cannabidiol (CBD) demonstrated antiarthritic effects independent of cannabinoid receptors. In addition to controlling inflammation, cannabinoids reduce pain by activating central and peripheral CB₁, peripheral CB₂ receptors and CBD-sensitive noncannabinoid receptor targets.

Summary

Cannabinoids might be a suitable treatment for RA, but it is important to target the right receptors in the right place. For clinical studies, we propose a combination of a CB₂ agonist to decrease cytokine production, a peripheral CB₁ antagonist to prevent detrimental CB₁ signaling and to support anti-inflammatory effects of CB₂ via activation of β_2 -adrenergic receptors and CBD to induce cannabinoid-receptor-independent anti-inflammatory effects.

Keywords

cannabidiol, cannabinoid, cytokines, rheumatoid arthritis, tetrahydrocannabinol

INTRODUCTION

Cannabis has been used for millennia as treatment for a wide variety of diseases such as depression, impotence, arthritis and kidney stones [1]. From the beginning of the 19th century until its complete ban in the United States in 1970, the psychopharmacological effects of cannabis were described in detail and phytocannabinoids were included in many pharmaceutical products. Significantly, the opioid sparing effect of cannabinoids was already known at the beginning of the 20th century and therefore, the plant was also used to treat opioid dependence [1].

Cannabis sativa contains over seventy different cannabinoids, with tetrahydrocannabinol (THC) and the nonpsychotropic cannabidiol (CBD) being the most prominent and best characterized [2]. Although THC mainly binds and activates the two cannabinoid receptors in humans, cannabinoid type 1 and 2 receptors (CB₁ and CB₂), CBD acts as a negative allosteric modulator at CB₁ and CB₂,

limiting the effects of THC, but it also binds and activates several transient receptor potential (TRP) ion channels, orphan G-protein coupled receptors (GPCRs) and it is involved in serotonergic neurotransmission [3,4,5,6,7,8]. Most effects of cannabis formulations are attributed to activation/modulation of CB₁ and CB₂. Although CB₁ is the predominant cannabinoid receptor in the central nervous system, CB₂ is distributed mainly in the periphery with high expression in immune cells [9,10]. In the central nervous system, cannabinoid

Poliklinik, Funktionsbereich & Hiller Forschungszentrum für Rheumatologie, University Hospital Duesseldorf, Duesseldorf, Germany

Correspondence to Dr Torsten Lowin, PhD, Hiller Forschungszentrum Rheumatologie, Life Science Center, Merowingerplatz1A, 1. Etage, 40225 Duesseldorf, Germany. Tel: +49 2118106150; fax: +49 2118106153; e-mail: torsten.lowin@med.uni-duesseldorf.de

Curr Opin Rheumatol 2019, 31:271–278

DOI:10.1097/BOR.0000000000000590

KEY POINTS

- CB₂ activation mediates anti-inflammatory effects in RA by decreasing immune cell migration and cytokine production.
- CB₁ activation promotes proinflammatory effects in immune cells, but reduces pain and depression.
- CB₁ antagonists provide anti-inflammatory effects by enhancing β₂-adrenergic signaling in arthritis.
- CBD is effective in reducing inflammation and pain and might enhance the efficacy of therapeutic drugs.
- A cannabinoid-based RA therapy would preferentially activate CB₂, inhibit CB₁ peripherally and include CBD to target noncannabinoid pathways.

receptors regulate neurotransmitter release and, depending on their concentration, modulate inhibitory and excitatory neurotransmission [11,12[■]]. Peripheral effects of cannabinoids include modulation of pain pathways, cytokine and immunoglobulin production, control of immune cell migration and control of glucose/energy supply [13–18,19[■],20].

The endogenous ligands for CB₁ and CB₂ are termed endocannabinoids with 2-arachidonylglycerol (2-AG) and anandamide (AEA) being the most prominent compounds [21,22]. Like phytocannabinoids, the action of endocannabinoids is not restricted to cannabinoid receptors but they also modulate TRP channels, gamma-aminobutyric acid receptors and several GPCRs [8,23,24]. Although the effects of exogenous cannabinoids are long lasting, endocannabinoids are rapidly (within minutes) inactivated enzymatically. Fatty acid amide hydrolase (FAAH) is the main degrading enzyme for AEA, while 2-AG is deactivated by monoacylglycerol lipase amongst others [25,26]. Significantly, AEA and 2-AG can also be metabolized by oxygenases, especially cyclooxygenase-2 (COX-2), yielding novel compounds with a pharmacology different from the parental endocannabinoids [27]. This mechanism might be especially important in (chronic) inflammatory situations where endocannabinoid metabolism might be diverted to COX-2, which is upregulated by proinflammatory cytokines [28]. In line with this, dual inhibitors for FAAH and COX-2 have been developed that synergize in increasing endocannabinoid levels [29].

Due to their anti-inflammatory effects, (endo-)cannabinoids have been considered as a potential therapy for the treatment of rheumatic diseases. However, at the moment there is no significant evidence for the efficacy of cannabinoid-based drugs

in the treatment of rheumatoid arthritis (RA), although it is included in the list of conditions eligible for receiving medical cannabis in Canada and several US states and the majority of patients with arthritis reported beneficial effects of the drug, for example less pain and an opioid-sparing effect [30,31[■],32]. Although human data are lacking, cannabinoids have been used in animal models of arthritis providing evidence for their clinical efficacy [33–36]. In this review, we discuss possible antiarthritic effects of cannabinoids and we hypothesize that cannabinoids might also help controlling RA-associated comorbidities (e.g. insulin resistance, hypertension) which is of special importance as they contribute to the increased mortality of RA patients [37].

DIRECT EFFECTS OF CANNABINOIDS ON THE IMMUNE SYSTEM: IT CAN GO EITHER WAY

RA is characterized by immune cell infiltration into the joint, augmented cytokine and chemokine production by resident fibroblasts and macrophages and cartilage destruction by matrix degrading enzymes [38,39]. Consequently, current RA therapeutics (including glucocorticoids) often interfere with cytokine production or signaling but are often associated with side effects related to infection or immune disturbances [40–42]. Therefore, reduction of proinflammatory cytokine production and signaling without an overt risk of infection would be a preferable treatment of RA. In vitro data and results from animal experiments suggest that cannabinoids might help with just that. Early studies in the 1970s already demonstrated that THC inhibits macrophage function [43,44]. In later experiments, it was demonstrated that CB₂ is the target receptor in macrophages to mediate anti-inflammatory effects. CB₂ reduces the activation of the NLRP3 inflammasome and subsequent IL-1β release, enhances clearance of apoptotic cells and generally decreases cellular activation in response to tumor necrosis factor (TNF) or lipopolysaccharide (LPS) [45–48]. Recently, the importance of CB₂ in RA development was emphasized by the finding that a loss-of function gene polymorphism increased the risk for disease 10 fold [49[■]]. In addition, inhibiting AEA degradation with subsequent increase in CB₂ activation reduced arthritis score and progression in a mouse model of mice with collagen-induced arthritis [36]. In contrast to CB₂, CB₁ induces proinflammatory effects by promoting macrophage M1 polarization, generation of reactive oxygen species, fibrosis and enhancing TLR4 signaling [50,51[■], 52,53]. This might be prevented by the use of CB₁

antagonists, but unfortunately, brain penetrable CB₁ antagonists, like rimonabant, demonstrated psychiatric side effects which led to discontinuation of the drug [54]. This was followed by the development of peripherally restricted CB₁ antagonists which might circumvent these central side effects [55].

Similar effects of cannabinoids have been demonstrated with cells from the adaptive immune system. In T cells, proliferation, differentiation and nuclear factor kappa-light-chain-enhancer of activated B cells activity were inhibited, while in B cells, migration, cellular energy supply and response to T-cell-independent antigens were modulated by CB₂ activation [56–59].

Although THC regulates several aspects of adaptive and innate immunity, an involvement of cannabinoid receptors is questionable as high concentrations of THC were necessary to elicit anti-inflammatory effects and effects were not shown to be inhibited by CB₁/CB₂ antagonists [60–63]. Although the mechanism of action is still unclear, CBD and its analogues demonstrated promising results in combating chronic inflammation in RA [33,35]. CBD induces regulatory T cells, activates and desensitizes several TRP ion channels, ligates peroxisome proliferator-activated receptor- γ and binds several other orphan GPCRs which might explain its complex pharmacology [4[■],7,64–66]. Like CBD, endocannabinoids engage several cellular targets other than cannabinoid receptors and some anti-inflammatory effects are attributed to activation of noncannabinoid related pathways [67].

When considering direct therapeutic effects of cannabinoids for the treatment of RA, it can be concluded that CB₂ agonists induce beneficial anti-inflammatory effects by downregulating pro-inflammatory cytokine production and reducing mobilization and migration of immune cells to sites of inflammation CB₁ agonists promote pro-inflammatory signaling and noncannabinoid, but endocannabinoid and phytocannabinoid (CBD)-sensitive receptors mediate anti-inflammatory effects.

In this respect, a combination of a CB₂ agonist and a peripherally restricted CB₁ antagonist together with CBD would hypothetically be an optimal combination to combat excessive inflammation and cytokine production in RA. Significantly, compounds with both, agonistic action at CB₂ and antagonistic action at CB₁, have already been described but not yet tested in models of chronic inflammation [68]. Similar therapeutic effects might be achieved by combining an endocannabinoid degradation inhibitor (preferentially a FAAH inhibitor) with a CB₁ antagonist.

INDIRECT ACTIONS OF CANNABINOIDS: TARGETING COMORBIDITIES IN RHEUMATOID ARTHRITIS

In addition to having direct effects on immune cells, cannabinoids might exert many additional effects on chronic inflammation via modulation of the sympathetic nervous system (SNS) and its neurotransmitters. It has already been demonstrated that anxiolytic and antidepressant effects or side effects of cannabinoids are dependent on the modulation of noradrenergic and serotonergic neurotransmission [12[■]]. In addition, the beneficial effects of CB₁ antagonism in a murine model of arthritis was dependent on intact β 2-adrenergic signaling and CB₁ was found to form dimers with β 2-adrenergic receptor [69,70].

Many RA patients suffer from comorbidities such as depression and cardiovascular events and it has been shown that hypertension and osteoporosis correlate with disease activity [71,72]. Decreased parasympathetic and an increase in sympathetic activity is one cause for the development of these conditions and, as they counteract neurotransmitter imbalances, cannabinoids might help in this respect [73–75]. In animal models, systemic activation of the SNS at the early phase of arthritis constitutes a proinflammatory signal as it mobilizes leukocytes, increases antigen uptake/presentation and provides energy to the immune system [76]. In addition, direct anti-inflammatory effects of sympathetic neurotransmitters on immune cells in the joint are blunted due to the repulsion of sympathetic nerve fibers from the site of inflammation [77]. In the later phase of the disease, compensatory, catecholamine-producing cells appear in the joint whose ablation aggravates experimental arthritis [78]. The question arises, how would cannabinoids interfere with the SNS in RA?

Cannabinoids modulate the outflow of sympathetic neurotransmitters centrally and peripherally through CB₁ and in the early phase of arthritis or to postpone clinical symptoms, reduction of SNS activity might be beneficial [12[■]]. Hypertension is a widespread problem in RA and it is often accompanied by insulin resistance and metabolic disturbances [79]. Activation of CB₁ reduces blood pressure and this effect was due to a reduction in noradrenergic tone [80]. Significantly, the consumption of cannabis is also associated with lower risk of metabolic syndrome and diabetes and might therefore prevent the RA-induced changes in metabolism [81]. This seems paradoxical as cannabis is known for its stimulating effect on food intake, but due to the partial agonism of THC, it also acts as antagonist for endocannabinoids [82,83]. In addition, CB₁ desensitizes and downregulates rather quickly resulting in functional antagonism even in the presence of agonist although

the rate and speed of this process is dependent on cell type and context [84]. Although CB₁ agonism might have some beneficial effects in the treatment of RA comorbidities, CB₁ antagonists are better suited to directly target inflammation in RA. Indeed, it has been shown that the anti-inflammatory effect of peripherally restricted CB₁ antagonists is dependent on intact adrenergic signaling in the spleen. CB₁ antagonism increases adrenergic β₂ signaling which reduces TNF production and arthritis severity [70]. In addition, metabolic changes due to low level systemic inflammation in RA might be diminished with CB₁ antagonists. In fact, this has been observed in the treatment of obesity and CB₁ antagonists improved insulin levels, lowered blood glucose and promoted a lean phenotype [82]. Another mechanism by which CB₁ antagonism might contribute to an anti-inflammatory phenotype in RA is by increasing local norepinephrine levels in joint and spleen that allow β-adrenergic signaling [70,85]. Norepinephrine has higher affinity for proinflammatory α-adrenergic receptors than for β-receptors resulting in mostly proinflammatory α-effects under low levels of norepinephrine [86]. As sympathetic nerves are repelled from the joint in the course of RA, norepinephrine levels are low serving an α-adrenergic dependent, proinflammatory role [77]. Like in the LPS-induced arthritis mouse model [70], use of a CB₁ antagonist might be able to increase intraarticular norepinephrine concentrations due to disinhibition at sympathetic nerve terminals recovering anti-inflammatory β-adrenergic effects. Although not tested experimentally, catecholamine-producing cells that appear in the course of arthritis might also be modulated by cannabinoids. The presence of the monoamine transporter VMAT2 suggests that catecholamines are stored in vesicles in tyrosine hydroxylase positive, catecholaminergic cells and their release might be under the control of CB₁ just like their neuronal counterparts [87]. If this is true, CB₁ activation would counteract catecholamine release and promote α-adrenergic, proinflammatory effects while CB₁ inhibition would be beneficial by increasing catecholamine levels, thereby promoting β-adrenergic effects.

CANNABINOIDS AND PAIN

The reduction of neuropathic and cancer pain are known indications for the use of cannabinoids and three cannabis-based drugs (Nabilone, Sativex and Marinol) are used clinically for this purpose [88,89]. Although there are only very limited clinical data on the effect of cannabinoids on arthritic pain, reports from patients show that a significant number uses cannabis to treat their symptoms [90]. In addition one study investigated the effects of sativex, a

combination of CBD and THC, in arthritis patients and this drug demonstrated a significant analgesic effect with mild adverse events like dizziness, but these did not lead to withdrawal from the study [91]. Although data from RA patients regarding cannabinoid effects are scarce, models of arthritic, inflammatory and neuropathic pain demonstrated a clear reduction of pain and allodynia when animals were treated with cannabinoids [92,93,94[■]]. In addition, data from human cohorts suggest at least limited efficacy of cannabis-based products on neuropathic pain [95]. Cannabinoids modulate pain by activating CB₁ and CB₂ receptors but some compounds also directly ligate the nociceptors TRP vanilloid 1 (TRPV1) and TRP Ankyrin 1 (TRPA1) [92,94[■],96]. Although CB₁ desensitizes nociceptors like the capsaicin receptor TRPV1 directly, CB₂ might reduce pain by inhibiting proinflammatory cytokine production and immune cell infiltrates that promote nociceptor function [97]. In addition, it has been demonstrated that CB₂ activation induces the peripheral release of β-endorphin, which might contribute to CB₂-induced analgesia [98,99]. Furthermore, CB₂ might functionally interact with peripheral μ-opioid receptors to induce antinociception and animals treated with a CB₂ agonist along with morphine showed additive analgesic effects [100]. Although CB₂-mediated analgesia is peripherally restricted, CB₁ reduces pain by peripheral and central mechanisms [101]. TRPV1 is a direct target of cannabinoids but it is also an ionotropic receptor for endocannabinoids [97,102,103]. AEA is able to desensitize TRPV1 and animals with complete FAAH inhibition demonstrate increased pain thresholds and TRPV1 antagonists or desensitizing agonists resulted in decreased proinflammatory cytokine production by macrophages [104–107]. Cannabinoids also support opioid-induced analgesia and it has been demonstrated that a combination of low dose THC with morphine prevents μ-opioid receptor desensitization and provides superior pain relief [89,108].

In addition to CB₁ and CB₂, other pathways are involved in cannabinoid-induced nociception. It has been shown that the nonpsychotropic phytocannabinoid CBD promotes analgesia by activating serotonin 5-HT_{1a} and TRPV1 receptors while AEA provides additional analgesic effects via peroxisome-proliferator activated receptor α [109[■],110].

CONCLUSION: WHAT WOULD BE A PREFERABLE CANNABINOID-BASED TREATMENT FOR RHEUMATOID ARTHRITIS?

CB₁ activation is accompanied by central, psychotropic side effects but also by peripheral,




FIGURE 1. Consequences of a peripherally restricted cannabinoid type 1 receptor antagonist combined with cannabidiol and a fatty acid amide hydrolase inhibitor on arthritic pain and inflammation. (1) Use of a peripherally restricted cannabinoid type 1 receptor antagonist increases local norepinephrine release and promotes anti-inflammatory β_2 -adrenergic signaling (indicated by the yield sign). (2) In addition, proinflammatory cannabinoid type 1 receptor signaling on immune cells is inhibited while anti-inflammatory effects are enhanced (indicated by the stop sign). (3) Fatty acid amide hydrolase inhibition increases systemic endocannabinoid levels. (4) This is accompanied by an increase in anti-inflammatory cannabinoid type 2 receptor signaling in macrophages, synovial fibroblasts, B cells and T cells. (5) Endocannabinoids but also cannabidiol desensitize and therefore inhibit the nociceptors TRP vanilloid 1 (TRPV1) and transient receptor potential Ankyrin 1 (TRPA1) which supports antiinflammation in immune cells but also inhibit pain transmission in sensory nerves. (6) Fatty acid amide hydrolase inhibition also increases central endocannabinoid levels. This allows for central, cannabinoid type 1 receptor signaling and inhibition of TRP channels resulting in pain relief and attenuation of depressive symptoms. TRP, transient receptor potential.

pro-inflammatory effects on immune cells that limit their clinical use [50,111]. In contrast, CB₁ activation provides pain relief and improves mood and one possibility to selectively activate mostly beneficial CB₁ pathways is by the means of FAAH inhibition. It has been shown that in contrast to THC, FAAH inhibition does not produce adverse effects but still promotes antidepressive behavior and analgesia [112–114]. Peripheral FAAH inhibition shares some of the negative effects of CB₁ activation in respect of metabolic changes and should therefore be avoided [115]. To prevent peripheral CB₁ activation a FAAH inhibitor might be combined with a CB₁ antagonist. With this combination, anti-

inflammatory effects would be augmented: FAAH inhibition elevates endocannabinoid levels in the brain that counteract depression and pain, while CB₁ antagonism in the periphery inhibits detrimental, pro-inflammatory CB₁ effects while boosting anti-inflammatory effects by increasing sympathetic nervous activity (Fig. 1). FAAH inhibition also enhances anti-inflammatory CB₂ signaling due to enhanced levels of AEA [36] (Fig. 1). In addition, CBD might be added to RA therapy, as it elicits antidepressive and anti-inflammatory effects independent of cannabinoid receptors and without psychotropic side effects. In arthritis animal models, CBD provided pain relief and reduced inflammatory

cell infiltrates into the joint [116]. Furthermore, CBD might boost the effect of antirheumatic drugs as it has been shown that it increases the uptake of chemotherapeutic compounds into cancer cells [7].

Acknowledgements

None.

Financial support and sponsorship

The work was supported by HILLER foundation, Monheim, Germany.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Pisanti S, Bifulco M. Medical cannabis: a plurimillennial history of an ever-green. *J Cell Physiol* 2018; doi: 10.1002/jcp.27725. [Epub ahead of print]
2. Peschel W, Politi M. (1)H NMR and HPLC/DAD for *Cannabis sativa* L. chemotype distinction, extract profiling and specification. *Talanta* 2015; 140:150–165.
3. Sales AJ, Crestani CC, Guimaraes FS, *et al.* Antidepressant-like effect induced by Cannabidiol is dependent on brain serotonin levels. *Prog Neuropsychopharmacol Biol Psychiatry* 2018; 86:255–261.

The study shows that the antidepressant effect of cannabidiol (CBD) is dependent on brain serotonin levels.

4. Laun AS, Shrader SH, Brown KJ, *et al.* GPR3, GPR6, and GPR12 as novel molecular targets: their biological functions and interaction with cannabidiol. *Acta Pharmacol Sin* 2018. [Epub ahead of print]

The study identified CBD as an antagonist at the orphan receptors GPR3, GPR6 and GPR12.

5. Iannotti FA, Hill CL, Leo A, *et al.* Nonpsychoactive plant cannabinoids, cannabidivarin (CBDV) and cannabidiol (CBD), activate and desensitize transient receptor potential vanilloid 1 (TRPV1) channels in vitro: potential for the treatment of neuronal hyperexcitability. *ACS Chem Neurosci* 2014; 5:1131–1141.
6. Navarro G, Reyes-Resina I, Rivas-Santisteban R, *et al.* Cannabidiol skews biased agonism at cannabinoid CB1 and CB2 receptors with smaller effect in CB1–CB2 heteroreceptor complexes. *Biochem Pharmacol* 2018; 157:148–158.

The study shows that functional selectivity of tetrahydrocannabinol and CP55940 is lost in cannabinoid type 1 receptor (CB₁)/cannabinoid type 2 receptor (CB₂) heterodimers. In addition, pretreatment with CBD resulted in enhanced functionality of the receptor complex.

7. Nabissi M, Morelli MB, Santoni M, *et al.* Triggering of the TRPV2 channel by cannabidiol sensitizes glioblastoma cells to cytotoxic chemotherapeutic agents. *Carcinogenesis* 2013; 34:48–57.
8. Kapur A, Zhao P, Sharir H, *et al.* Atypical responsiveness of the orphan receptor GPR55 to cannabinoid ligands. *J Biol Chem* 2009; 284:29817–29827.
9. Katona I. Endocannabinoid receptors: CNS localization of the CB(1) cannabinoid receptor. *Curr Top Behav Neurosci* 2009; 1:65–86.
10. Galiegue S, Mary S, Marchand J, *et al.* Expression of central and peripheral cannabinoid receptors in human immune tissues and leukocyte subpopulations. *Eur J Biochem* 1995; 232:54–61.
11. Grzeda E, Schlicker E, Toczek M, *et al.* CB1 receptor activation in the rat paraventricular nucleus induces bi-directional cardiovascular effects via modification of glutamatergic and GABAergic neurotransmission. *Naunyn Schmiedeberg's Arch Pharmacol* 2017; 390:25–35.
12. Mendiguren A, Aostri E, Pineda J. Regulation of noradrenergic and serotonergic systems by cannabinoids: relevance to cannabinoid-induced effects. *Life Sci* 2018; 192:115–127.

The review shows the influence of CB₁ on adrenergic and serotonergic neurotransmission.

13. Cencioni MT, Chirurciu V, Catanzaro G, *et al.* Anandamide suppresses proliferation and cytokine release from primary human T-lymphocytes mainly via CB2 receptors. *PLoS One* 2010; 5:e8688.

14. Guabiraba R, Russo RC, Coelho AM, *et al.* Blockade of cannabinoid receptors reduces inflammation, leukocyte accumulation and neovascularization in a model of sponge-induced inflammatory angiogenesis. *Inflamm Res* 2013; 62:811–821.
15. Chirurciu V, Cencioni MT, Bisicchia E, *et al.* Distinct modulation of human myeloid and plasmacytoid dendritic cells by anandamide in multiple sclerosis. *Ann Neurol* 2013; 73:626–636.
16. Jorda MA, Verbakel SE, Valk PJ, *et al.* Hematopoietic cells expressing the peripheral cannabinoid receptor migrate in response to the endocannabinoid 2-arachidonoylglycerol. *Blood* 2002; 99:2786–2793.
17. Agudelo M, Newton C, Widen R, *et al.* Cannabinoid receptor 2 (CB2) mediates immunoglobulin class switching from IgM to IgE in cultures of murine-purified B lymphocytes. *J Neuroimmune Pharmacol* 2008; 3:35–42.
18. Basu S, Ray A, Dittel BN. Cannabinoid receptor 2 is critical for the homing and retention of marginal zone B lineage cells and for efficient T-independent immune responses. *J Immunol* 2011; 187:5720–5732.
19. Chan LN, Chen Z, Braas D, *et al.* Metabolic gatekeeper function of B-lymphoid transcription factors. *Nature* 2017; 542:479–483.

The study demonstrates a new role of CB₂ in the utilization of energy in B cells and identified CB₂ as therapeutic target to treat B-cell malignancies.

20. Romero TR, Resende LC, Guzzo LS, *et al.* CB1 and CB2 cannabinoid receptor agonists induce peripheral antinociception by activation of the endogenous noradrenergic system. *Anesth Analg* 2013; 116:463–472.
21. Vogel Z, Barg J, Levy R, *et al.* Anandamide, a brain endogenous compound, interacts specifically with cannabinoid receptors and inhibits adenylate cyclase. *J Neurochem* 1993; 61:352–355.
22. Di M, De VPL, Sugiura T, *et al.* Potential biosynthetic connections between the two cannabimimetic eicosanoids, anandamide and 2-arachidonoylglycerol, in mouse neuroblastoma cells. *Biochem Biophys Res Commun* 1996; 227:281–288.
23. Ross RA. Anandamide and vanilloid TRPV1 receptors. *Br J Pharmacol* 2003; 140:790–801.
24. Musella A, Fresegna D, Rizzo FR, *et al.* A novel crosstalk within the endocannabinoid system controls GABA transmission in the striatum. *Sci Rep* 2017; 7:7363.
25. Ueda N, Puffenbarger RA, Yamamoto S, *et al.* The fatty acid amide hydrolase (FAAH). *Chem Phys Lipids* 2000; 108:107–121.
26. Dinh TP, Carpenter D, Leslie FM, *et al.* Brain monoglyceride lipase participating in endocannabinoid inactivation. *Proc Natl Acad Sci U S A* 2002; 99:10819–10824.
27. Urquhart P, Nicolaou A, Woodward DF. Endocannabinoids and their oxygenation by cyclo-oxygenases, lipoxygenases and other oxygenases. *Biochim Biophys Acta* 2015; 1851:366–376.
28. Alhouayek M, Muccioli GG. COX-2-derived endocannabinoid metabolites as novel inflammatory mediators. *Trends Pharmacol Sci* 2014; 35:284–292.
29. Scarpelli R, Sasso O, Piomelli D. A double whammy: targeting both fatty acid amide hydrolase (FAAH) and cyclooxygenase (COX) to treat pain and inflammation. *ChemMedChem* 2016; 11:1242–1251.
30. Walitt B, Klose P, Fitzcharles MA, *et al.* Cannabinoids for fibromyalgia. *Cochrane Database Syst Rev* 2016; 7:CD011694.
31. Katz-Talmor D, Katz I, Porat-Katz BS, *et al.* Cannabinoids for the treatment of rheumatic diseases – where do we stand? *Nat Rev Rheumatol* 2018; 14:488–498.

The review gives an overview about the anti-inflammatory effects of cannabis and discusses clinical and preclinical data that support cannabis-based therapies in rheumatic disease.

32. Ko GD, Bober SL, Mindra S, *et al.* Medical cannabis – the Canadian perspective. *J Pain Res* 2016; 9:735–744.
33. Malfait AM, Gallily R, Sumariwalla PF, *et al.* The nonpsychoactive cannabis constituent cannabidiol is an oral antiarthritic therapeutic in murine collagen-induced arthritis. *Proc Natl Acad Sci U S A* 2000; 97:9561–9566.
34. Sofia RD, Nalepa SD, Harakal JJ, *et al.* Antiedema and analgesic properties of delta9-tetrahydrocannabinol (THC). *J Pharmacol Exp Ther* 1973; 186:646–655.
35. Sumariwalla PF, Gallily R, Tchilibon S, *et al.* A novel synthetic, nonpsychoactive cannabinoid acid (HU-320) with antiinflammatory properties in murine collagen-induced arthritis. *Arthritis Rheum* 2004; 50:985–998.
36. Kinsey SG, Naidu PS, Cravatt BF, *et al.* Fatty acid amide hydrolase blockade attenuates the development of collagen-induced arthritis and related thermal hyperalgesia in mice. *Pharmacol Biochem Behav* 2011; 99:718–725.
37. Marrie RA, Walld R, Bolton JM, *et al.* Psychiatric comorbidity increases mortality in immune-mediated inflammatory diseases. *Gen Hosp Psychiatry* 2018; 53:65–72.
38. Bartok B, Firestein GS. Fibroblast-like synoviocytes: key effector cells in rheumatoid arthritis. *Immunol Rev* 2010; 233:233–255.
39. Smolen JS, Aletaha D, McInnes IB. Rheumatoid arthritis. *Lancet* 2016; 388:2023–2038.
40. Dhillon S. Tocilizumab: a review in rheumatoid arthritis. *Drugs* 2017; 77:1987–2001.
41. Geng Z, Yu Y, Hu S, *et al.* Tocilizumab and the risk of respiratory adverse events in patients with rheumatoid arthritis: a systematic review and meta-analysis of randomised controlled trials. *Clin Exp Rheumatol* 2018. [Epub ahead of print]

42. Caporali R, Crepaldi G, Codullo V, *et al.* 20 years of experience with tumour necrosis factor inhibitors: what have we learned? *Rheumatology (Oxford)* 2018; 57:vi5–vii10.
43. Gaul CC, Mellors A. Delta-9-tetrahydrocannabinol and decreased macrophage migration inhibition activity. *Res Commun Chem Pathol Pharmacol* 1975; 10:559–564.
44. Raz A, Goldman R. Effect of hashish compounds on mouse peritoneal macrophages. *Lab Invest* 1976; 34:69–76.
45. Persidsky Y, Fan S, Dykstra H, *et al.* Activation of cannabinoid type 2 receptors (CB2) diminish inflammatory responses in macrophages and brain endothelium. *J Neuroimmune Pharmacol* 2015; 10:302–308.
46. Jiang L, Chen Y, Huang X, *et al.* Selective activation of CB2 receptor improves efferocytosis in cultured macrophages. *Life Sci* 2016; 161:10–18.
47. Ke P, Shao BZ, Xu ZQ, *et al.* Activation of cannabinoid receptor 2 ameliorates DSS-induced colitis through inhibiting NLRP3 inflammasome in macrophages. *PLoS One* 2016; 11:e0155076.
48. Gui H, Liu X, Liu LR, *et al.* Activation of cannabinoid receptor 2 attenuates synovitis and joint destruction in collagen-induced arthritis. *Immunobiology* 2015; 220:817–822.
49. Ismail M, Khawaja G. Study of cannabinoid receptor 2 Q63R gene polymorphism in Lebanese patients with rheumatoid arthritis. *Clin Rheumatol* 2018; 37:2933–2938.
- The study shows the influence of a gain-of function mutation of CB₂ on the development of rheumatoid arthritis.
50. Lecru L, Desterke C, Grassin-Delyle S, *et al.* Cannabinoid receptor 1 is a major mediator of renal fibrosis. *Kidney Int* 2015; 88:72–84.
51. Ruiz dA, Mancini IG, Srivastava RK, *et al.* Adipocyte cannabinoid receptor CB1 regulates energy homeostasis and alternatively activated macrophages. *J Clin Invest* 2017; 127:4148–4162.
- The study shows a strong influence of white adipocyte-expressed CB₁ on whole body energy expenditure and metabolic parameters.
52. Duncan M, Galic MA, Wang A, *et al.* Cannabinoid 1 receptors are critical for the innate immune response to TLR4 stimulation. *Am J Physiol Regul Integr Comp Physiol* 2013; 305:R224–R231.
53. Han KH, Lim S, Ryu J, *et al.* CB1 and CB2 cannabinoid receptors differentially regulate the production of reactive oxygen species by macrophages. *Cardiovasc Res* 2009; 84:378–386.
54. Moreira FA, Crippa JA. The psychiatric side-effects of rimonabant. *Rev Bras Psiquiatr* 2009; 31:145–153.
55. Silvestri C, Di Marzo V V. Second generation CB1 receptor blockers and other inhibitors of peripheral endocannabinoid overactivity and the rationale of their use against metabolic disorders. *Expert Opin Investig Drugs* 2012; 21:1309–1322.
56. Kozela E, Haj C, Hanus L, *et al.* HU-446 and HU-465, derivatives of the nonpsychoactive cannabinoid cannabidiol, decrease the activation of encephalitogenic T cells. *Chem Biol Drug Des* 2016; 87:143–153.
57. Robinson RH, Meissler JJ, Fan X, *et al.* A CB2-selective cannabinoid suppresses T-cell activities and increases Tregs and IL-10. *J Neuroimmune Pharmacol* 2015; 10:318–332.
58. Kong W, Li H, Tuma RF, *et al.* Selective CB2 receptor activation ameliorates EAE by reducing Th17 differentiation and immune cell accumulation in the CNS. *Cell Immunol* 2014; 287:1–17.
59. Singh UP, Singh NP, Singh B, *et al.* Cannabinoid receptor-2 (CB2) agonist ameliorates colitis in IL-10(–/–) mice by attenuating the activation of T cells and promoting their apoptosis. *Toxicol Appl Pharmacol* 2012; 258:256–267.
60. Klein TW, Newton CA, Widen R, *et al.* The effect of delta-9-tetrahydrocannabinol and 11-hydroxy-delta-9-tetrahydrocannabinol on T-lymphocyte and B-lymphocyte mitogen responses. *J Immunopharmacol* 1985; 7:451–466.
61. Schatz AR, Koh WS, Kaminski NE. Delta 9-tetrahydrocannabinol selectively inhibits T-cell dependent humoral immune responses through direct inhibition of accessory T-cell function. *Immunopharmacology* 1993; 26:129–137.
62. Bhargava HN, House RV, Thorat SN, *et al.* Cellular immune function in mice tolerant to or abstinent from l-trans-delta 9-tetrahydrocannabinol. *Pharmacology* 1996; 52:271–282.
63. Zhu W, Friedman H, Klein TW. Delta-9-tetrahydrocannabinol induces apoptosis in macrophages and lymphocytes: involvement of Bcl-2 and caspase-1. *J Pharmacol Exp Ther* 1998; 286:1103–1109.
64. Dhital S, Stokes JV, Park N, *et al.* Cannabidiol (CBD) induces functional Tregs in response to low-level T cell activation. *Cell Immunol* 2017; 312:25–34.
65. De PL, Orlando P, Moriello AS, *et al.* Cannabinoid actions at TRPV channels: effects on TRPV3 and TRPV4 and their potential relevance to gastrointestinal inflammation. *Acta Physiol (Oxf)* 2012; 204:255–266.
66. O'Sullivan SE, Kendall DA. Cannabinoid activation of peroxisome proliferator-activated receptors: potential for modulation of inflammatory disease. *Immunobiology* 2010; 215:611–616.
67. Lowin T, Apitz M, Anders S, *et al.* Anti-inflammatory effects of N-acylethanolamines in rheumatoid arthritis synovial cells are mediated by TRPV1 and TRPA1 in a COX-2 dependent manner. *Arthritis Res Ther* 2015; 17:321.
68. Dhopeshwarkar A, Murataeva N, Makriyannis A, *et al.* Two janus cannabinoids that are both CB2 agonists and CB1 antagonists. *J Pharmacol Exp Ther* 2017; 360:300–311.
69. Hudson BD, Hebert TE, Kelly ME. Physical and functional interaction between CB1 cannabinoid receptors and beta2-adrenoceptors. *Br J Pharmacol* 2010; 160:627–642.
70. Mnich SJ, Hiebsch RR, Huff RM, *et al.* Anti-inflammatory properties of CB1-receptor antagonist involves beta2 adrenoceptors. *J Pharmacol Exp Ther* 2010; 333:445–453.
71. Klodzinski L, Wislowska M. Comorbidities in rheumatic arthritis. *Reumatologia* 2018; 56:228–233.
72. Dougados M, Soubrier M, Antunez A, *et al.* Prevalence of comorbidities in rheumatoid arthritis and evaluation of their monitoring: results of an international, cross-sectional study (COMORA). *Ann Rheum Dis* 2014; 73:62–68.
73. del RA, Wolff C, Wildmann J, *et al.* Disrupted brain-immune system-joint communication during experimental arthritis. *Arthritis Rheum* 2008; 58:3090–3099.
74. Adlan AM, Paton JF, Lip GY, *et al.* Increased sympathetic nerve activity and reduced cardiac baroreflex sensitivity in rheumatoid arthritis. *J Physiol* 2017; 595:967–981.
75. Godfrey KEM, Gardner AC, Kwon S, *et al.* Differences in excitatory and inhibitory neurotransmitter levels between depressed patients and healthy controls: a systematic review and meta-analysis. *J Psychiatr Res* 2018; 105:33–44.
76. Pongratz G, Straub RH. The sympathetic nervous system in inflammation. *Arthritis Res Ther* 2014; 16:504.
77. Fassold A, Falk W, Anders S, *et al.* Soluble neuropilin-2, a nerve repellent receptor, is increased in rheumatoid arthritis synovium and aggravates sympathetic fiber repulsion and arthritis. *Arthritis Rheum* 2009; 60:2892–2901.
78. Capellino S, Weber K, Gelder M, *et al.* First appearance and location of catecholaminergic cells during experimental arthritis and elimination by chemical sympathectomy. *Arthritis Rheum* 2012; 64:1110–1118.
79. Medina G, Vera-Lastra O, Peralta-Amaro AL, *et al.* Metabolic syndrome, autoimmunity and rheumatic diseases. *Pharmacol Res* 2018; 133:277–288.
80. Malinowska B, Toczek M, Pedzinska-Betkik A, *et al.* Cannabinoids in arterial, pulmonary and portal hypertension – mechanisms of action and potential therapeutic significance. *Br J Pharmacol* 2018. [Epub ahead of print]
81. Sidney S. Marijuana use and type 2 diabetes mellitus: a review. *Curr Diab Rep* 2016; 16:117.
82. Salamone JD, McLaughlin PJ, Sink K, *et al.* Cannabinoid CB1 receptor inverse agonists and neutral antagonists: effects on food intake, food-reinforced behavior and food aversions. *Physiol Behav* 2007; 91:383–388.
83. Lu HC, Mackie K. An introduction to the endogenous cannabinoid system. *Biol Psychiatry* 2016; 79:516–525.
84. Schlosburg JE, Blankman JL, Long JZ, *et al.* Chronic monoacylglycerol lipase blockade causes functional antagonism of the endocannabinoid system. *Nat Neurosci* 2010; 13:1113–1119.
85. Lowin T, Straub RH. Cannabinoid-based drugs targeting CB1 and TRPV1, the sympathetic nervous system, and arthritis. *Arthritis Res Ther* 2015; 17:226.
86. Pongratz G, Straub RH. The sympathetic nervous response in inflammation. *Arthritis Res Ther* 2014; 16:504.
87. Capellino S, Cosentino M, Wolff C, *et al.* Catecholamine-producing cells in the synovial tissue during arthritis: modulation of sympathetic neurotransmitters as new therapeutic target. *Ann Rheum Dis* 2010; 69:1853–1860.
88. Russo EB, Guy GW, Robson PJ. Cannabis, pain, and sleep: lessons from therapeutic clinical trials of Sativex, a cannabis-based medicine. *Chem Biodivers* 2007; 4:1729–1743.
89. Mucke M, Phillips T, Radbruch L, *et al.* Cannabis-based medicines for chronic neuropathic pain in adults. *Cochrane Database Syst Rev* 2018; 3:CD012182.
90. Ware MA, Adams H, Guy GW. The medicinal use of cannabis in the UK: results of a nationwide survey. *Int J Clin Pract* 2005; 59:291–295.
91. Blake DR, Robson P, Ho M, *et al.* Preliminary assessment of the efficacy, tolerability and safety of a cannabis-based medicine (Sativex) in the treatment of pain caused by rheumatoid arthritis. *Rheumatology (Oxford)* 2006; 45:50–52.
92. Cox ML, Haller VL, Welch SP. The antinociceptive effect of Delta-9-tetrahydrocannabinol in the arthritic rat involves the CB(2) cannabinoid receptor. *Eur J Pharmacol* 2007; 570:50–56.
93. Petrovski Z, Kovacs G, Tomboly C, *et al.* The effects of peptide and lipid endocannabinoids on arthritic pain at the spinal level. *Anesth Analg* 2012; 114:1346–1352.
94. Klinger-Gratz PP, Ralvenius WT, Neumann E, *et al.* Acetaminophen relieves inflammatory pain through CB1 cannabinoid receptors in the rostral ventromedial medulla. *J Neurosci* 2018; 38:322–334.
- The study identified CB₁ as one cellular target of acetaminophen to counteract pain generation in the ventromedial medulla.
95. Nugent SM, Morasco BJ, O'Neil ME, *et al.* The effects of cannabis among adults with chronic pain and an overview of general harms: a systematic review. *Ann Intern Med* 2017; 167:319–331.
96. De PL, Vellani V, Schiano-Moriello A, *et al.* Plant-derived cannabinoids modulate the activity of transient receptor potential channels of ankyrin type-1 and melastatin type-8. *J Pharmacol Exp Ther* 2008; 325:1007–1015.
97. Jeske NA, Patwardhan AM, Gamper N, *et al.* Cannabinoid WIN 55,212-2 regulates TRPV1 phosphorylation in sensory neurons. *J Biol Chem* 2006; 281:32879–32890.

98. Ibrahim MM, Porreca F, Lai J, *et al.* CB2 cannabinoid receptor activation produces antinociception by stimulating peripheral release of endogenous opioids. *Proc Natl Acad Sci U S A* 2005; 102:3093–3098.
99. Gao F, Zhang LH, Su TF, *et al.* Signaling mechanism of cannabinoid receptor-2 activation-induced beta-endorphin release. *Mol Neurobiol* 2016; 53:3616–3625.
100. Yuill MB, Hale DE, Guindon J, *et al.* Antinociceptive interactions between opioids and a cannabinoid receptor 2 agonist in inflammatory pain. *Mol Pain* 2017; 13:1744806917728227.
101. Fox A, Kesingland A, Gentry C, *et al.* The role of central and peripheral Cannabinoid1 receptors in the antihyperalgesic activity of cannabinoids in a model of neuropathic pain. *Pain* 2001; 92:91–100.
102. Toth A, Blumberg PM, Boczan J. Anandamide and the vanilloid receptor (TRPV1). *Vitam Horm* 2009; 81:389–419.
103. Wang ZY, McDowell T, Wang P, *et al.* Activation of CB1 inhibits NGF-induced sensitization of TRPV1 in adult mouse afferent neurons. *Neuroscience* 2014; 277:679–689.
104. Tsuji F, Murai M, Oki K, *et al.* Transient receptor potential vanilloid 1 agonists as candidates for anti-inflammatory and immunomodulatory agents. *Eur J Pharmacol* 2010; 627:332–339.
105. Ninomiya Y, Tanuma SI, Tsukimoto M. Differences in the effects of four TRPV1 channel antagonists on lipopolysaccharide-induced cytokine production and COX-2 expression in murine macrophages. *Biochem Biophys Res Commun* 2017; 484:668–674.
106. Starowicz K, Makuch W, Korostynski M, *et al.* Full inhibition of spinal FAAH leads to TRPV1-mediated analgesic effects in neuropathic rats and possible lipoxygenase-mediated remodeling of anandamide metabolism. *PLoS One* 2013; 8:e60040.
107. Lizanecz E, Bagi Z, Pasztor ET, *et al.* Phosphorylation-dependent desensitization by anandamide of vanilloid receptor-1 (TRPV1) function in rat skeletal muscle arterioles and in Chinese hamster ovary cells expressing TRPV1. *Mol Pharmacol* 2006; 69:1015–1023.
108. Cichewicz DL, McCarthy EA. Antinociceptive synergy between delta(9)-tetrahydrocannabinol and opioids after oral administration. *J Pharmacol Exp Ther* 2003; 304:1010–1015.
109. De Gregorio D, McLaughlin RJ, Posa L, *et al.* Cannabidiol modulates serotonergic transmission and reverses both allodynia and anxiety-like behavior in a model of neuropathic pain. *Pain* 2019; 160:136–150.
- The study shows that CBD acts on central TRP vanilloid 1 and 5-HT1a receptors to regulate analgesia and anxiety.
110. Kaczocha M, Rebecchi MJ, Ralph BP, *et al.* Inhibition of fatty acid binding proteins elevates brain anandamide levels and produces analgesia. *PLoS One* 2014; 9:e94200.
111. Tanda G, Goldberg SR. Cannabinoids: reward, dependence, and underlying neurochemical mechanisms – a review of recent preclinical data. *Psychopharmacology (Berl)* 2003; 169:115–134.
112. Kangas BD, Leonard MZ, Shukla VG, *et al.* Comparisons of Delta9-tetrahydrocannabinol and anandamide on a battery of cognition-related behavior in nonhuman primates. *J Pharmacol Exp Ther* 2016; 357:125–133.
113. Wang Y, Zhang X. FAAH inhibition produces antidepressant-like effects of mice to acute stress via synaptic long-term depression. *Behav Brain Res* 2017; 324:138–145.
114. Ahn K, Johnson DS, Mileni M, *et al.* Discovery and characterization of a highly selective FAAH inhibitor that reduces inflammatory pain. *Chem Biol* 2009; 16:411–420.
115. Vaitheesvaran B, Yang L, Hartil K, *et al.* Peripheral effects of FAAH deficiency on fuel and energy homeostasis: role of dysregulated lysine acetylation. *PLoS One* 2012; 7:e33717.
116. Hammell DC, Zhang LP, Ma F, *et al.* Transdermal cannabidiol reduces inflammation and pain-related behaviours in a rat model of arthritis. *Eur J Pain* 2016; 20:936–948.